



QUESTIONS MAY 2010

CPD questionnaires must be completed online via www.cpdjournals.org.za. After submission you can check the answers and print your certificate. Questions may be answered up to 6 months after publication of each issue.

1. True (A) or false (B) – click on the correct answer:
The presence of multiple metaphyseal fractures is highly suggestive of child abuse.
2. True (A) or false (B) – click on the correct answer:
Umbilical cord blood is a useful source of stem cells for regenerative transplantation.
3. True (A) or false (B) – click on the correct answer:
Fortunately, the costs of establishing umbilical cord blood banks are negligible.
4. True (A) or false (B) – click on the correct answer:
Ritonavir is a protease inhibitor used in the treatment of advanced HIV infection.
5. True (A) or false (B) – click on the correct answer:
Ritonavir inhibits cytochrome P450-CYP3A4 iso-enzyme activity in the liver.
6. True (A) or false (B) – click on the correct answer:
Adrenal suppression with ritonavir and concomitant corticosteroid, fluticasone, has been described in children.
7. True (A) or false (B) – click on the correct answer:
A recent study has shown that budesonide can also produce adrenal suppression in children when used with ritonavir.
8. True (A) or false (B) – click on the correct answer:
Hookahs are also known as water pipes.
9. True (A) or false (B) – click on the correct answer:
Hookah tobacco smokers have minimal health risks compared with cigarette smokers.
10. True (A) or false (B) – click on the correct answer:
Only tobacco is used by hookah smokers.
11. True (A) or false (B) – click on the correct answer:
In South Africa hookah smoking is only done by adults.
12. True (A) or false (B) – click on the correct answer:
Young hookah smokers claim that they mainly smoke for health reasons.
13. True (A) or false (B) – click on the correct answer:
The World Health Organization estimated that there were >800 000 malaria deaths in Africa in 2006.
14. True (A) or false (B) – click on the correct answer:
Most deaths from malaria in Africa are of children.
15. True (A) or false (B) – click on the correct answer:
The Eastern Cape has become a high-risk region for malarial transmission.
16. True (A) or false (B) – click on the correct answer:
***Plasmodium falciparum* accounts for the vast majority of malarial infections diagnosed in South Africa.**
17. True (A) or false (B) – click on the correct answer:
Imported malaria in South Africa is largely confined to affluent recreational travellers.
18. True (A) or false (B) – click on the correct answer:
Delivery by caesarean section (CS) is one of the most common major operative procedures performed worldwide.
19. True (A) or false (B) – click on the correct answer:
Morphine and pethidine are not transferred to breastmilk.
20. True (A) or false (B) – click on the correct answer:
A review showed that women undergoing CS under regional analgesia who had local anaesthetic infiltration had a reduced need for postoperative opioids.

INSTRUCTIONS

1. Read the journal. All the answers will be found there.
2. Go to www.cpdjournals.org.za to answer the questions.

Accreditation number: MDB001/015/01/2009 (Clinical)