CPD questionnaires must be completed online via www.cpdjournals.org.za. After submission you can check the answers and print your certificate. Questions may be answered up to 6 months after publication of each issue.

- 1. True (A) or false (B) click on the correct answer: Most of the 5 million HIV-positive South Africans are older than the reproductive age group.
- 2. True (A) or false (B) click on the correct answer: **HIV** infection results in enhanced fertility.
- 3. True (A) or false (B) click on the correct answer:

 Data from a rape centre showed that in 84.5% of cases <15
 years old, the perpetrator was known to the survivor.
- 4. True (A) or false (B) click on the correct answer: Raped males were younger and presented with more genital and anal injuries.
- 5. True (A) or false (B) click on the correct answer: Warfarin-induced skin necrosis (WISN) is a rare complication of warfarin therapy.
- 6. True (A) or false (B) click on the correct answer: Classic histological features of WISN include full-thickness epidermal necrosis and thrombosed vessels in the dermis.
- 7. True (A) or false (B) click on the correct answer: WISN is fortunately associated with low morbidity and is never fatal.
- 8. True (A) or false (B) click on the correct answer: Most cases of WISN occur after treatment for venous thrombo-embolism (VTE).
- 9. True (A) or false (B) click on the correct answer: WISN never occurs in patients with cardiac indications for warfarin therapy.
- 10. True (A) or false (B) click on the correct answer:

 Women are affected with WISN more frequently than men.
- 11. True (A) or false (B) click on the correct answer: Mycobacterium tuberculosis may present clinically as deepvein thrombosis.

- 12. True (A) or false (B) click on the correct answer: **Because of their low CD4 counts, patients with HIV have no risk of developing VTE.**
- 13. True (A) or false (B) click on the correct answer: Coal combustion and goldmining are sources of mercury poisoning.
- 14. Which of the following is NOT correct? Other causes of mercury exposure include:
- a) Dental fillings containing mercury
- b) Consumption of contaminated fish
- c) Breakage of mercury-containing devices, e.g. thermometers
- d) Excessive consumption of very old whisky.
- 15. True (A) or false (B) click on the correct answer: **Methylmercury is toxic even at low concentrations.**
- 16. True (A) or false (B) click on the correct answer: Gold amalgamation is used by small-scale miners because it is efficient in extracting gold and is inexpensive.
- 17. True (A) or false (B) click on the correct answer: When mercury levels in the urine are high, patients may experience symptoms such as fever, tremors, mood swings and respiratory symptoms.
- 18. True (A) or false (B) click on the correct answer: Cryptococcal meningitis (CM) is now the most common cause of adult meningitis in southern Africa.
- 19. True (A) or false (B) click on the correct answer: Since CM is easily treatable, it is never fatal.
- 20. True (A) or false (B) click on the correct answer: The recommended treatment of HIV-associated CM in South Africa is fluconazole consolidation for all patients, following initial amphotericin B therapy.

INSTRUCTIONS

- $1. \ Read \ the journal.$ All the answers will be found there.
- 2. Go to www.cpdjournals.org.za to answer the questions.

Accreditation number: MDB001/015/01/2009 (Clinical)

TIOZZAIR